


Percentage of general primary care and primary care subspecialty physicians who are certified in palliative medicine, 2000

Range 0.00 – 0.97% Mean 0.33%



- | | | | | | |
|--|---|--|---|---|--|
| <p>A Maine
Montana
New Hampshire
New Mexico
Vermont</p> | <p>B Arizona
Florida
Iowa
Kansas
Michigan
Ohio
Oklahoma
Utah
Wyoming</p> | <p>C California
Colorado
Connecticut
Delaware
District of Columbia
Hawaii
Illinois
Kentucky
Louisiana
Maryland
Minnesota
Mississippi
Missouri
New York
Oregon</p> | <p>Rhode Island
South Dakota
Tennessee
Texas
Washington
West Virginia
Wisconsin</p> | <p>D Alabama
Arkansas
Indiana
Massachusetts
Nevada
New Jersey
North Carolina
North Dakota
Pennsylvania</p> | <p>E Alaska
Georgia
Idaho
Nebraska
North Dakota
South Carolina
Virginia</p> |
|--|---|--|---|---|--|

Source: American Board of Hospice and Palliative Medicine; www.abhpm.org. American Medical Association. *Physician Characteristics and Distribution in the US*, 2002–2003 Edition (Chicago: American Medical Association Press, 2002).