


Strength of state pain policies, 2001

Range -2 - +9



- | | | | | |
|--|---|--|---|--|
| <p>A Alabama
Florida
Kansas
Nebraska
North Carolina
Utah
Washington</p> | <p>B Kentucky
Maine
Maryland
Minnesota
Missouri
Nevada
Pennsylvania
Rhode Island
West Virginia</p> | <p>C Arizona
California
Colorado
Montana
New Hampshire
New Mexico
Oklahoma
Oregon
South Carolina
South Dakota
Tennessee
Wyoming</p> | <p>D Arkansas
District of Columbia
Georgia
Iowa
Massachusetts
Michigan
Mississippi
New Jersey
North Dakota
Ohio
Texas
Vermont
Virginia</p> | <p>E Alaska
Connecticut
Delaware
Hawaii
Idaho
Illinois
Indiana
Louisiana
New York
Wisconsin</p> |
|--|---|--|---|--|

Source: Based on *Achieving Balance in Federal & State Pain Policy*, dated July 2000, and updated in the *Annual Review of State Pain Policies 2000* (published February 2001) and *Annual Review of State Pain Policies 2001* (published February 2002) from the Pain and Policy Studies Group of the University of Wisconsin's Comprehensive Cancer Center, Madison, Wisconsin. More recent information may be available at the group's Web site: www.medsch.wisc.edu/painpolicy.